

# WITT MORPHISMS



NR 2966

PRZEMYSŁAW KOPROWSKI

# WITT MORPHISMS

Wydawnictwo Uniwersytetu Śląskiego



Katowice 2012

Redaktor serii: Matematyka

**Tomasz Dłotko**

Recenzent

**Grzegorz Banaszak**

Publikacja będzie dostępna — po wyczerpaniu nakładu — w wersji internetowej:

Śląska Biblioteka Cyfrowa

[www.sbc.org.pl](http://www.sbc.org.pl)

# CONTENTS

<b>Foreword</b> . . . . .	7
<b>1. Preliminaries</b> . . . . .	11
1.1. Witt functor . . . . .	11
1.2. Orderings and valuations . . . . .	15
1.3. Knebusch-Milnor exact sequence . . . . .	20
1.4. Introduction to real curves . . . . .	23
<b>2. Local quadratic extensions</b> . . . . .	29
2.1. Transfer maps . . . . .	32
2.2. Scharlau's norm principle . . . . .	37
Appendix: Unitary extensions of arbitrary degree . . . . .	39
<b>3. Injectivity of Witt functor of ring normalization</b> . . . . .	45
3.1. Preparatory lemmas . . . . .	45
3.2. Locally unitary quadratic extensions . . . . .	49
3.3. Relations to the Picard functor . . . . .	53
Appendix: Curve desingularization . . . . .	56
<b>4. Splitting the Knebusch-Milnor exact sequence</b> . . . . .	61
4.1. Preparatory lemmas . . . . .	61
4.2. Direct sum theorem . . . . .	63
4.3. Splitting theorem . . . . .	68
4.4. Further splitting . . . . .	69
<b>5. Witt equivalence of real rings</b> . . . . .	73
5.1. Witt equivalence of real fields . . . . .	73
5.2. Witt equivalence of real holomorphy rings . . . . .	75
Appendix: Quaternion-symbol equivalence of geometric rings . . . . .	80
<b>Bibliography</b> . . . . .	85
<b>Commonly used symbols</b> . . . . .	89
<b>Index</b> . . . . .	91
Streszczenie . . . . .	95
Zusammenfassung . . . . .	95

# FOREWORD

A bilinear form defines an orthogonal geometry on a given linear space or projective module. Once we omit degenerate forms (roughly speaking, these are forms allowing a non-zero vector to be orthogonal to the entire space), the remaining ones may be classified by the relation of similarity (the term is explained in Section 1.1). This leads us to the notion of the Witt ring, which is an algebraic structure consisting of all the similarity classes of finitely generated projective modules over a given base ring. In a sense, Witt ring describes all the possible orthogonal geometries over the ring/field in question. The leading theme of this book is to study morphisms between Witt rings. In geometric terms, this can be viewed as analyzing to what extent the orthogonal geometries defined over one ring may be transcribed to another ring. For example, knowing the criteria for existence of an isomorphism between Witt rings, one may verify whether two rings/fields admit the same set of orthogonal geometries. If this is the case we say that the two rings are *Witt equivalent*. This problem has been intensively researched in previous years. The complete criteria are known for fields with small square class groups (see [9]) and global fields (see e.g. [44, 53, 52]). The author of this book found criteria for Witt equivalence of function fields and rings of regular function on real algebraic curves (summary of these results can be found in the appendix to Chapter 5). The criterion of Witt equivalence of real function fields has been recently generalized by N. Grenier-Boley and D. Hoffmann to arbitrary real fields with  $u$ -invariants not exceeding 2. We present their result in Section 5.1. In the second section of the last chapter we apply the ideas used earlier for rings of regular function on real curves to extend the result of Grenier-Boley and Hoffmann and obtain a necessary condition (see Theorem 5.15) for Witt equivalence of real holomorphy rings. This generalizes our earlier result obtained in [27].

Chapter 4 copes with a splitting of the Knebusch-Milnor exact sequence. A classical theorem due to M. Knebusch and J. Milnor (see Theorem 1.40) asserts that the Witt ring  $WR$  of a Dedekind domain  $R$  injects into the Witt ring of its field of fractions  $K$  and the image of this embedding coincides with the kernel of a map from the Witt ring of the field to the co-product of Witt rings of all the localizations of the base ring. In a nutshell, this says that the structure of

orthogonal geometries over  $K$  is at least as rich as the structure over the underlying Dedekind domain. Surprisingly, not much is known about the splitting of the above-mentioned injection  $WR \rightarrow WK$ . The problem was solved in the case of algebraic integers by P. Shastri in [50] and for real geometric rings by the author in [26] and [28]. Chapter 4 summarizes the results of these two papers. The main result of this chapter (see Theorem 4.3) asserts that, if  $R$  is the ring of regular functions on a smooth real curve, then  $WR$  is a direct summand of the Witt ring of the field of fractions of  $R$  (the field of rational functions on this curve). Consequently, the Knebusch-Milnor exact sequence slices into and is patched by two split exact sequences (c.f. Theorem 4.17). Moreover, if the curve in question is semi-algebraically compact and semi-algebraically connected, then the Witt ring of the ring of polynomial function is in turn a direct summand of  $WR$  as shown in Theorem 4.19.

On the other hand, it is natural to reckon that starting from a ring with a complex Witt ring (i.e. with a rich structure of orthogonal geometries) and appending roots of (quadratic) polynomials one can successively kill elements of the Witt ring. Therefore, it is expected that the natural morphism of Witt rings corresponding to an algebraic (resp.: real, quadratic or integral) closure of a field/ring should *not* be injective. A classical example: start from the field  $\mathbb{Q}$  of rationals, let  $R$  denote its real closure and  $R[\sqrt{-1}]$  be the algebraic closure. The Witt ring of  $\mathbb{Q}$  has quite a complex structure (as additive group it is a direct sum of infinitely many nontrivial terms, see [33, Chapter VI, Section 4]), while  $WR$  is isomorphic to the ring of integers and  $WR[\sqrt{-1}]$  consists of just two elements. Thus, the injections  $\mathbb{Q} \rightarrow R \rightarrow R[\sqrt{-1}]$  correspond to the maps  $W\mathbb{Q} \rightarrow WR \rightarrow WR[\sqrt{-1}]$ , both having strongly nontrivial kernels. In Chapter 3 we concentrate on an analogy of this phenomenon in the case of the integral closure of a ring. For example, we show (see Theorem 3.15) that if  $P$  is seminormal but not quadratically closed and  $R$  is the integral closure of  $P$ , then the natural morphism  $WP \rightarrow WR$  is not injective. This problem has also a natural interpretation in terms of the Picard functor. This connection is studied in Section 3.3. We close this chapter showing how to apply these results in the case of curve desingularization.

The Witt functor of a ring extension is also the subject of the second chapter. Here, however, we consider a quadratic extension of a local ring. We develop a generalization of Scharlau's transfer and prove an analogy of Scharlau's norm principle. This allows us to construct examples of ring extensions where both rings have the same field of fractions but the corresponding Witt morphism is not surjective (hence there are classes of forms over the bigger ring not coming from the smaller one).

The new contributions in this book include: entire Chapters 2 and 3 and Section 5.2 where the main new results are Theorems: 2.17, 2.19, 2.24, 3.11, 3.15, 3.24, 5.15 and Proposition 2.35. The results of Chapter 4 appeared earlier in

[26, 28]. The proofs presented here are only slightly improved and unified to better fit together. The results gathered in the appendix to Chapter 5 were published in [24, 25, 27, 29]. All the theorems of Chapter 1 are classical and well known. The intend of the author was to make the book as self-contained as possible, to save the reader from the need to refer to any external sources while reading. Hence the opening chapter contains a basic introduction to the theory of bilinear forms, valuations and orderings, serving as a handy reference for the following chapters. The presentation of the first chapter is necessarily brief and most of the proofs are omitted.

A short remark on the notational convention. We tend to use letters  $P$  and  $R$  to denote rings. While the latter is widely used and self-explaining (as the first letter of the word “ring”), the use of  $P$  requires some justification. We often need to compare two rings and so  $R$  alone is not enough. The letter  $P$  is close enough to  $R$  so that we can write “ $P$  is a subring of  $R$ ” with the inclusion relation preserving the natural alphabetical order. Secondly,  $P$  is the first letter of the Polish word “pierścień” (which means ring), hence we have  $P$  for “pierścień” and  $R$  for “ring”. The rest of notation used in this book is standard and agrees with broadly accepted conventions. For reader’s convenience, we include the list of commonly used symbols on page 89.



# INDEX

- adjoint homomorphism, 12
- algebraic function field, 24
- Artin's theorem, 17, 74
- Artin-Schreier theorem, 16
  
- Baer-Krull theorem, 18, 24, 76
- bilinear form, 11, 34, 35, 37, 49
  - anisotropic, 74
  - binary, 37, 39, 42, 77
  - diagonal, 13, 34, 37, 40
  - hyperbolic, *see* hyperbolic form
  - isotropic, 26
  - non-degenerate, 12, 34, 38, 41
  - on free module, 13, 34
  - on vector space, 12, 21, 22
  - over Noetherian ring, 14, 47, 57
  - unary, 23, 35, 42, 77
- bilinear module, *see* bilinear form
- blow-up, 59
  
- component (of curve), 25, 26, 61–65, 80, 82
  - homeomorphic to circle, 25, 26, 81
  - separation, 25, 65
- component (of space of real places), 79
- conductor, 32, 33, 35, 37, 39, 48, 50, 53–55, 57
- curve desingularization, 36, 56–59
- curve normalization, *see* curve desingularization
  
- Dedekind domain, 20, 20, 22, 23, 27, 48, 69, 75–78
- definite
  - form, 25
  - function, 25, 62, 64, 65
- determinant, 30
  - of bilinear form, 13, 14, 35–38, 42
- discrete valuation, 17, 18, 20
- discriminant, 13, 70, 77
  
- Euclidean topology, 24
- exceptional divisor, 59
  
- field
  - complete, 17, 63
  - real, *see* formally real field
  - real closed, *see* real closed field
- field of fractions, 16, 46, 47, 50, 52, 53
  - of Dedekind ring, 20, 22, 23, 48, 49, 76
  - of local ring, 30
- form
  - bilinear, *see* bilinear form
  - quadratic, *see* quadratic form
- formally real field, 16, 19, 61, 63, 73, 74, 76–78, 80, 82
- free module, 13, 29, 32–34, 40, 46, 63
  
- genus, 58, 82
  
- Harrison theorem, 74, 75
- Harrison topology, 20, 74, 75
- Harrison's criterion, *see* Harrison's theorem
- Harrison's isomorphism, 73, 78
- Harrison's theorem, 73
- Hasse principle, 26
- hyperbolic form, 13, 14, 21, 22, 29, 31, 32, 36–39, 41, 42, 45, 47, 48, 57, 75, 76

- indefinite
  - form, 26
- integral closure, *see* ring normalization
- interval (on real curve), 26, 62, 63, 66, 81
- interval function, 26, 62–66, 71
  - uniqueness, 26, 62
- isometric forms, *see* isometry
- isometry, 12, 26, 63, 71, 72
  - class, 13
- isotropic vector
  - strongly, 14, 47
  - weakly, 14
- Jacobson's radical, 30
- Knebusch-Milnor sequence, 20–23, 27, 48, 49, 61–72, 82
  - for real curve, 27
- Knebusch-Milnor theorem, *see* Knebusch-Milnor sequence
- lattice, 21–23
- line bundle, 14, 51
- local ring, 13, 17, 22, 29, 30, 34, 35, 37–39, 42
  - regular, *see* regular local ring
- localization
  - commutes with intersection, 50
  - of ideal, 46, 49
  - of ring, 48, 52
- lying-over theorem, 56
- Mayer-Vietoris sequence, 53, 54
- metabolic form, *see* hyperbolic form
- module
  - bilinear, *see* bilinear form
  - free, *see* free module
  - projective, *see* projective module
  - torsion-free, *see* torsion-free module
- nilpotent element, 19
- nilradical, 19, 23
  - of Witt ring, 19, 23, 76, 77, 79
- Noetherian ring, 14, 20, 29, 30, 35, 47–52, 56
- normalization
  - of curve, *see* curve desingularization
  - of ring, *see* ring normalization
- ordering, 15, 16, 75–77
  - and field's characteristic, 16
  - compatible with point, 24–26, 65, 79, 81
  - compatible with valuation, 17, 18, 76, 78
  - from signature, 19
  - on field of fractions, 16
  - on residue field, 16, 76, 78
  - support, *see* support of ordering
  - unique, 16, 24, 78
- orientation (of curve), 26
- orthogonal
  - basis, 13
  - geometry, 15, 58
  - sum, 12, 15, 32, 35
  - vectors, 12, 15, 21, 41
- Pfister form, 13, 19, 39, 75
- Pfister's local-global principle, 19, 75–77
- Picard
  - functor, 45, 51, 53, 56
  - group, 51, 55, 57
- principal ideal, 14, 47, 49, 50, 55, 57
- principal ideal domain, 17
- projective module, 11–13, 21, 32, 46, 48, 49
  - of rank 1, *see* line bundle
  - over Dedekind domain, 20
  - over Noetherian ring, 14
  - over semi-local ring, 13
- proper transform, 59
- quadratic form, 12, 25, 32, 76
  - associated to bilinear form, 12
  - binary, 63, 71, 73, 75
  - definite, *see* definite form
  - hyperbolic, *see* hyperbolic form
  - unary, 63, 65, 67, 70, 71, 76
- quaternion-symbol equivalence, 80–83
- tame, 78

- real algebraic curve, 23–27, 36, 57, 61,  
     69, 72, 77, 80–82  
     singular, 36, 56, 59  
 real closed field, 16, 24, 75, 78, 80  
     of constants, 24, 36, 56, 57, 59, 61,  
     63, 80, 82  
 real field, *see* formally real field  
 real holomorphy ring, 73, 75–78  
 real point, 24–27, 61–66, 69, 71, 72, 79,  
     81, 82  
     nonsingular, 36  
     singular, 36  
 real valuation, 17, 18, 78, 80  
 reduced Witt ring, 77  
 regular local ring, 30, 31  
 regular prime, 51  
 residue field, 16, 24, 36, 78  
     of local ring, 29, 30, 40, 52  
     of valuation, 17, 18, 22, 51, 76, 78  
 residue homomorphism  
     first, 20  
     second, 20, 23, 27  
 ring  
     Dedekind, *see* Dedekind domain  
     local, *see* local ring  
     Noetherian, *see* Noetherian ring  
     normalization, 32, 35, 45, 48, 49,  
     51, 53, 56  
     of regular functions, 27, 36, 58, 59,  
     61–70, 77, 80, 82  
     semi-local, *see* semi-local ring  
     seminormal, *see* seminormal ring  
 ring extension, 15, 32, 45, 53  
     integral, 56  
     local quadratic, 29, 31, 32, 40, 52  
     locally unitary, 45, 49, 50–52, 55–57  
     non-unitary, 29, 38  
     quadratic, 48, 49  
     subintegral, 51  
     unitary, 29, 39  
 ring normalization, 48  
  
 Scharlau's norm principle, 37–39  
 Scharlau's theorem, *see* Scharlau's norm  
     principle  
 Scharlau's transfer, *see* transfer  
 semi-algebraically connected component,  
     *see* component (of curve)  
 semi-local ring, 13, 29, 40  
 seminormal ring, 45, 52, 53  
 sign at a point, 25, 26, 62, 64, 66, 71, 72  
 signature, 19, 25, 76, 77  
     preserving map, 74  
     total, *see* total signature  
 similar forms, 14, 37  
 space of orderings, 16, 74, 75, 79  
     topology, *see* Harrison topology  
 strong isomorphism (of Witt rings), 74  
 strong topology, *see* Euclidean topology  
 support of ordering, 15, 16  
  
 tensor product  
     of bilinear modules, 12, 13, 15  
     of line bundles, 51  
 theorem  
     Artin's, *see* Artin's theorem  
     Artin-Schreier, *see* Artin-Schreier  
     theorem  
     Baer-Krull, *see* Baer-Krull theorem  
     Harrison, *see* Harrison theorem  
     Knebusch-Milnor, *see*  
     Knebusch-Milnor sequence  
     lying-over, *see* lying-over theorem  
     Scharlau's, *see* Scharlau's norm  
     principle  
     Traverso-Swan, *see* Traverso-Swan  
     theorem  
     Witt, *see* Witt theorem  
 torsion element, 19  
     in Picard group, 53  
     in Witt group, 19, 74, 77  
 torsion subgroup, 19  
     of Witt ring, 19, 76  
 torsion-free module, 20, 21  
 total signature, 19, 75  
 totally positive, 17, 62–65, 67, 77, 79  
 transfer, 31–38, 40, 42  
     proper, 34, 35, 36, 41, 42  
 Traverso-Swan theorem, 52  
  
 $u$ -invariant, 73–75

- uniformizer, 17, 20, 22, 23, 27, 76
- valuation, 17, 18
  - compatible with ordering, *see* ordering compatible with valuation
  - discrete, *see* discrete valuation
  - real, *see* real valuation
- valuation ring, 17, 21, 24, 36, 51, 69
  - residually real, 17, 76, 78, 80
  - trivial, 17
- vector space, 12, 21, 29
  - over  $\mathbb{F}_2$ , 74
- Witt equivalence, 73–83
- Witt functor, 11, 15, 31
  - of curve desingularization, 56
  - of natural injection, 29, 32
  - of non-unitary extension, 39
  - of ring normalization, 35, 45, 48, 49, 51, 53, 57, 59
- Witt group, 15, 36, 41, 57, 61, 63, 68, 69
  - of field, 32, 67, 70, 72
  - of local ring, 34
- Witt ring, 15, 32, 63, 64, 73, 75, 77
  - of curve, 27
  - of Dedekind ring, 61, 64, 76
  - of field, 63
  - of holomorphy ring, 75, 79
  - reduced, *see* reduced Witt ring
- Witt theorem, 26, 63, 71, 72, 75
- $\mathcal{X}$ -equivalence, 74, 75
  - tame, 75, 78
- Zariski topology, 19

Przemysław Koprowski

## MORFIZMY PIERŚCIENI WITTA

### S t r e s z c z e n i e

Rozprawa *Witt morphisms* omawia właściwości funktora Witt'a na kategorii pierścieni przemiennych z jedyneką. Książka składa się z pięciu rozdziałów, z których pierwszy ma charakter wprowadzający do tematyki funktora Witt'a. W rozdziale tym są zdefiniowane kluczowe pojęcia niezbędne do rozumienia dalszych części pracy i przywołane standardowe wyniki używane w kolejnych rozdziałach. Główną część pracy stanowią rozdziały 2–5. Rozdział drugi omawia problematykę zachowania funktora Witt'a na rozszerzeniach unitarnych (a w szczególności na kwadratowych rozszerzeniach unitarnych) pierścieni lokalnych. Rozdział ten zawiera między innymi uogólnienie techniki transferowej Scharlau'ego na przypadek rozszerzeń niewolnych. Rozdział trzeci wykorzystuje wyniki rozdziału poprzedniego do badania zachowania funktora Witt'a normalizacji dziedzin wymiaru jeden. W rozdziale tym w szczególności poruszana jest kwestia (nie)injektywności funktora Witt'a normalizacji. Rozdział czwarty poświęcony jest tematyce rozszczepialności ciągu dokładnego Knebuscha–Milnora dla pierścieni geometrycznych. Ostatni, piąty, rozdział rozprawy dotyczy równoważności Witt'a rzeczywistych ciał i pierścieni, czyli istnienia izomorfizmu między pierścieniami Witt'a dwóch struktur algebraicznych.

Przemysław Koprowski

## WITTMORPHISMEN

### Z u s a m m e n f a s s u n g

Das Buch *Witt morphisms* befasst sich mit den Eigenschaften des Wittfunktors in der Kategorie der kommutativen Ringe mit Eins. Das Buch hat fünf Kapitel. Das erste gibt eine Einführung in die Terminologie und die klassischen Resultate, die für die weiteren Kapitel notwendig sind. Die Hauptresultate des Buches sind in den Kapiteln 2–5 enthalten. Das zweite Kapitel diskutiert das Verhalten des Wittfunktors unter unitären Erweiterungen (insbesondere unter quadratischen unitären Erweiterungen) lokaler Ringe. Neben anderen Themen enthält dieses Kapitel eine Verallgemeinerung von Scharlaus Transferprinzip für nicht-freie Erweiterungen. Gegenstand des dritten Kapitels ist das Verhalten des Wittfunktors unter Normalisierung. Eines der Hauptthemen ist die Nicht-Injektivität des Wittfunktors unter Normalisierung. Das vierte Kapitel beschäftigt sich mit dem Problem der Zerfällung der Knebusch–Milnor exakten Folge für geometrische Ringe. Das letzte, fünfte Kapitel behandelt die Theorie der Witt-Äquivalenz formal reeller Ringe und Körper.

Redaktor  
Barbara Todos-Burny

Redaktor techniczny  
Barbara Arenhövel

Projektant okładki, skład i łamanie  
Przemysław Koprowski

Copyright © 2012 by  
Wydawnictwo Uniwersytetu Śląskiego  
Wszelkie prawa zastrzeżone

**ISSN 0208-6336**  
**ISBN 978-83-226-2113-4**

Wydawca  
**Wydawnictwo Uniwersytetu Śląskiego**  
**ul. Bankowa 12B, 40-007 Katowice**  
[www.wydawnictwo.us.edu.pl](http://www.wydawnictwo.us.edu.pl)  
e-mail: [wydawus@us.edu.pl](mailto:wydawus@us.edu.pl)

---

Wydanie I. Ark. druk. 6,0. Ark. wyd. 7,0.  
Papier offset. kl. III, 90 g      Cena 8 zł (+ VAT)

---

Druk i oprawa: PPHU TOTEM s.c.  
M. Rejnowski, J. Zamiara  
ul. Jacewska 89, 88-100 Inowrocław